

ANALYSIS OF RUMEN DEGRADATION CHARACTERISTICS OF FORAGE CRUDE PROTEIN IN GOAT

Andi Ikhsan WIJAYA , Ismartoyo ISMARTOYO  , and Asmuddin NATSIR 

Department of Nutrition and Animal Feed, Faculty of Animal Husbandry, Hasanuddin University, Perintis Kemerdekaan Road Km. 10, Makassar 902445, Indonesia

✉ Email: ismartoyo@gmail.com

➤ Supporting Information

ABSTRACT: The quality of feed given to ruminants can be determined from the degradation of nutrient content in the rumen. This study aimed to determine the pattern of forage degradation and the characteristics degradation of crude protein in the rumen using the *in sacco* method. The study used 4 fistulae kacang goats with an average body weight of 14.57 kg. The forage used consisted of R1: elephant grass (*Pennisetum purpureum*), R2: mini elephant grass (*Pennisetum purpureum* cv. *Mott*), R3: guinea grass (*Panicum maximum*), and R4: signal grass (*Bracharia decumbens*). The nylon bag is made of polyester measuring 8x4 cm with a porosity of 40µm. Feed samples were put into the rumen and incubated for 0, 4, 8, 12, 24, 48, and 72 hours. The parameters measured were consumption, patterns, and forage degradation characteristics by calculating the values of a, b, c, a+b, lag time, and ED. Determination characteristics of feed degradation in the rumen by *in sacco* method will be analyzed. The results showed that the characteristics of crude protein degradation had significant differences in fraction values a, b, a+b, and lag time ($P < 0.05$), while c and ED did not have significant differences ($P > 0.05$). In conclusion the crude protein of the degradation characteristics in the rumen were: elephant grass (a: 9.88%, b: 64.37%, and c: 0.06/h⁻¹), mini elephant grass (a: 16.50%, b: 45.24%, and c: 0.05/h⁻¹), guinea grass (a: 7.42%, b: 68.24%, and c: 0.05/h⁻¹), and signal grass (a: 6.79%, b: 56.19%, and c: 0.07/h⁻¹). So, grass can provide sufficient protein for microbial growth in the rumen for ruminants.

Keywords: Crude protein, Degradability, Forage, *in sacco*, Rumen.

RESEARCH ARTICLE
 PII: S222877012200033-13
 Received: December 11, 2023
 Revised: May 26, 2023
 Accepted: May 27, 2023

INTRODUCTION

Forage is the main source of energy and basic needs for ruminants, in general term (Minson, 1990). Feed costs can reach 50% - 70% of the production cost of a ruminant livestock business (Bozic et al., 2012). The forage that is generally given to ruminants is derived from the grass or gramineae group. Currently the main obstacle in increasing livestock production in tropical countries is the availability and quality of feed ingredients.

Ruminant farms in Indonesia generally use grass as the main feed which can be found nearby, for example elephant grass, mini elephant grass, guinea grass, and signal grass. The four grasses can be developed in tropical climates and have good nutrition to be used as feed for ruminants.

Feed is generally assessed for its quality, one of which is based on the protein content contained therein and aspects of degradation in the rumen. Provision of protein from forage to ruminants needs to pay attention to aspects of degradation in the rumen for microbial needs and those that escape microbial degradation (by-pass) as a source of protein to be utilized by the host (Puastuti et al., 2014). Protein degradation in the rumen is necessary to provide the N source needed for microbial growth (Mutsvangwa et al., 2016). Nichols et al. (2022) added that the contribution of microbial protein in the rumen plays an important role in sustaining N requirements in ruminants.

The *in sacco* method is a method for measuring the degradation value of a feed ingredient in the rumen (Mahrez and Ørskov, 1977; Guadayo et al., 2019). The degradation value of the sample will be measured with a nylon bag containing the sample and will be incubated in the rumen of fistula cattle at certain time intervals (Ørskov, 2000). It is important to know the quality of the protein content in elephant grass, mini elephant grass, guinea grass, and signal grass. It is necessary to conduct research to analyze the patterns and characteristics of feed degradation by using forage sources in livestock *in sacco*.

MATERIALS AND METHODS

Ethical approval

The experimental procedure for *in sacco* degradation of forage feed in live animals complies with the principles of animal welfare and was approved by the Health Research Ethics Committee, Hasanuddin University (Approved Number: 645/UN4.6.4.5.31/PP36/2021, Protocol UH21090601) prior to this study held.

Experimental animals and diets

This research was conducted from September 2021 to February 2022 at Hasanuddin University, Makassar, Indonesia. This study used the Latin Square Design method 4x4 which used 4 goats with fistulas of the male kacang variety (*Capra aegragus hircus*) aged ± 2 years. The average weight of the goats used as test animals was 14.57 ± 1.219 Kg. During the observation, the animals were given the same feed consisting of elephant grass, mini elephant grass, guinea grass, signal grass, and rice bran each of the goat's dry matter requirement, namely 3% of initial body weight (BB) given twice a day every morning and evening. Drinking water is provided *ad libitum*.

Experimental design

Observation of feed degradation characteristics was carried out using the *in sacco* method, using 8x4 cm nylon bags with 40 μ m porosity. The feed ingredients tested were R1: elephant grass (*Pennisetum purpureum*), R2: mini elephant grass (*Pennisetum purpureum* cv. *Mott*), R3: guinea grass (*Panicum maximum*), and R4: signal grass (*Bracharia decumbens*) which were harvested 40 days after the last pruning. The grass will be dried in an oven (60°C) for 24 hours and ground to a size of ± 2 mm then tested for proximate analysis (AOAC, 1995) and analysis of fiber components (Van Soest, 1994) the results can be seen in table 1. Each sample of feed ingredient 3 grams was put in a nylon bag and then incubated in the rumen with an incubation period of 0, 4, 8, 12, 24, 48, and 72 hours. This study used the Latin Square Design 4x4 randomization method which consisted of four observation periods using four goats and four different forages. The incubated bag was put into the oven at 60°C for 48 hours. During this observation, each animal was given the same feed consisting of the 4 tested feed ingredients with the addition of 20% rice bran.

Calculation of crude protein content in feed before and after incubation using the Kjeldahl method (AOAC, 2001) at the Feed Chemistry Laboratory, Faculty of Animal Husbandry, Hasanuddin University. The loss of crude protein in samples from nylon bags during each incubation period is assumed to be a protein that has been successfully degraded in the rumen used to calculate the degradation of crude protein feed according to the type of grass and the length of the incubation period. Determination of Crude Protein (PK) which is degraded in the sample can be known by multiplying the percentage of PK content in the initial sample. The formula for calculating the percentage loss of sample PK is presented as follows: % PK Loss = (%Initial PK x Initial Sample Weight) – (%Final PK x Final Sample Weight)

Furthermore, the crude protein lost during the incubation period is used to measure the value of Y by calculating the values a, b, c and a+b which are entered into the exponential equation according to Ørskov and McDonald (1979) as follows:

$$Y = a + b(1 - e^{-ct})$$

Where:

Y= Feed degradation by rumen microbes at time t (incubation time); a= soluble fraction; b= Potentially degraded fraction; c= Potential fraction degradation rate (b); a + b= Total potential degradation, including material that escapes the codend without degradation; t₀= Incubation time at 0 hours

The effectiveness of degradation is measured by the equation according to Orksov and McDonald (1979): $ED = a + [(b \times c)/(c + k)]$, assuming the feed flow rate (k) is 0.05/h (Srakaew et al., 2021). Effectively degraded crude protein is assumed to be Rumen Degradable Protein (RDP). Rumen Undegradable Protein (RUP) from each sample is calculated by the following equation: $RUP = 100\% - RDP$ (Terefe et al. 2022). Determination of the degradation curve and feed characteristics in the rumen *in sacco* will be analyzed using the Neway program (Ismartoyo, 2011). Data analysis uses One-Way Analysis of Variance (ANOVA), and if there are differences, then continue with Duncan's test. Statistical data used Software Package for Social Sciences (SPSS) Version 25. Then used Microsoft Excel 2010 software to see the degradation curve.

RESULTS AND DISCUSSION

Feed consumption

Consume fresh ingredients in livestock is at an average of 78.52 g/Kg BW/day. According to McDonald et al. (2002) factors that influence consumption in livestock are the characteristics of feed ingredients, environment, and livestock conditions. Han et al. (2019) added that palatability greatly affects the consumption of feed in livestock. The dry matter intake in this study was around 460.59 g/day or 61.68 g/Kg^{0.75}/day. These results are lower than those obtained by Mbuthia and Gachuri (2003), who obtained a value of 79 g/Kg^{0.75}/day, but higher than the results of Manaye et al. (2009) which was only 57.9 g/Kg^{0.75}/day and Han et al. (2019) who obtained a value of 369.9 g/day. The normal range of dry-matter intake per kilogram of metabolic bodyweight in goats is in the range of 34–104 g/Kg (Decandia et al., 2007). The voluntary intake of crude protein in goat poo is 56.33 g/day, or 7.55 g/Kg^{0.75}/day. These findings are similar to those obtained by Sultana et al. (2012), whose crude protein intake for goats ranged from 45.2 to 75.7 g/day, and Baete and Aregheore (2011), whose crude protein intake for grass was 7 g/Kg^{0.75}/day.

Crude protein degradation

Table 1 shows that the incubation period of 4, 8 and 12 hours of the four grasses had no significant difference (P<0.05). Whereas at 24, 48, and 72 hours incubation showed significant differences (P<0.05) from the various feeds

be negative (Ørksov et al., 1980; Odedire et al., 2013).

Table 2 - Consumption of livestock voluntary feed

Goat	Intake as feed basis (g/day)	DM Intake (g/day)	CP Intake (g/day)
Goat-1	1245.14±43.29	501.29±17.43	61.31±2.13
Goat-2	1154.67±93.54	464.87±37.66	56.85±4.61
Goat-3	1049.97±76.75	422.72±30.90	51.70±3.78
Goat-4	1126.33±41.40	453.46±16.67	55.46±2.04
Average	1144.03	460.59	56.33

Description: Feed consists of 20% elephant grass + 20% mini elephant grass + 20% guinea grass + 20% signal grass + 20% rice bran

Degradation effectiveness

The effectiveness of degradation (ED) is the result of the accumulation of feed degradation characteristics such as easily degradable fraction (a), slowly degraded fraction (b), and degradation rate of b fraction (c). EDCP on the four feeds showed results that were not significantly different (P>0.05). The effectiveness of crude protein degradation in the rumen of the four treatments showed a higher EDCP value at R1, namely 40.29% ± 2.47 although according to statistical tests it does not show the difference. This can be seen from the NH₃ level which is a product of feed protein content that is degraded by proteolytic enzymes in the rumen (Sari et al., 2021). In study of Mushandri (2022), which measured the concentration of NH₃ from the same four types of feed, the NH₃ level in elephant grass was the highest, 4.64 mM compared to other grasses. The effectiveness of crude protein degradation in elephant grass obtained a value of 43.20% ± 2.47 which is different from the results obtained from previous studies, namely 34.53% (da Silva et al., 2021) and 53.9% (Katongole et al., 2021). Mini elephant grass has a degradation effectiveness value of 37.88% ± 4.81, which is different from the results obtained by other researchers, namely 46.81%/d (da Silva et al., 2021), and 64% (Orsoletta et al., 2017). For Guinea grass, the degradation effectiveness value was obtained as much as 40.33% ± 2.72, this is different from other studies which obtained results of 34.18% (Ogunwole et al., 2011) and 46.67% (Bonelli et al., 2013). Meanwhile, based on the calculation of the effectiveness of degradation, Signal Grass obtained a value of 39.32% ± 5.54, this value is higher than the results obtained by Lana et al. (2007) namely 16.4% and Terefe et al. (2022) namely 27.86%. These different results were obtained because differences in the nutrient content of the samples could be caused by lighting/climate (Ballare et al., 1997), leaf/stem ratio (Lemaire et al., 2020), processing with fermentation (Ferreira et al., 2014), the addition of other materials and differences in results can also be caused by differences in the methods used.

Table 3- The average percentage of CP degradation at each incubation period

Incubation Period (Hours)	Elephant Grass (%)	Mini Elephant Grass (%)	Guinea Grass (%)	Signal Grass (%)
0	6.80 ± 2.85	4.05±1.65	6.32±1.73	2.68±0.91
4	19.58±1.70	12.75±1.79	14.04±3.22	12.50±3.68
8	27.81±1.34	23.87±0.50	27.96±5.81	26.82±4.98
12	36.47±5.57	31.07±1.67	35.60±5.24	34.78±6.24
24	54.87±3.55 ^b	42.08±0.75 ^a	46.86±2.82 ^a	47.34±8.79 ^a
48	68.99±4.86 ^b	60.53±1.41 ^a	67.57±2.80 ^b	59.68±6.91 ^a
72	71.47±3.99 ^b	62.13±1.88 ^a	70.80±1.72 ^b	62.00±6.31 ^a

a,b,c,d: Means in the same row with different superscripts differ significantly (P<0.05)

Table 4 - Characteristics of crude protein degradation

Parameters	R1	R2	R3	R4
A (%)	9.88±0.38 ^b	16.50±0.67 ^a	7.42±1.09 ^c	6.79±0.35 ^d
B(%)	64.37±4.85 ^{ab}	45.24±11.37 ^c	68.24±2.301 ^a	56.19±5.183 ^b
A+B (%)	74.25±4.85 ^a	61.74±11.37 ^b	75.66±2.31 ^a	62.98±5.17 ^b
C (/H)	0.06±0.012	0.05±0.006	0.05±0.008	0.07±0.016
Lt (h ⁻¹)	1.2±0.84 ^b	4.2±0.08 ^a	1.3±1.02 ^b	1.2±1.07 ^b
Ed (%)	43.20±2.47	37.88±4.51	40.33±2.72	39.32±5.54
Rup (%)	56.80±2.47	62.12±4.51	59.67±2.72	60.68±5.54

R1: elephant grass (*pennisetum purpureum*), R2: mini elephant grass (*pennisetum purpureum cv. Mott*), R3: guinea grass (*panicum maximum*), R4: signal grass (*brachiaria decumbens*), a: soluble fraction, b: potential degradation fraction, a+b: total potential degradation, c: degradation rate of fraction b, lt: lag time, ed: degradation effectiveness, rup: rumen undegradable protein. Different superscripts in the same row show significant differences (p<0.05)




Figure 1 - Crude protein degradation curve Elephant grass (*Pennisetum purpureum*), Mini elephant grass (*Pennisetum purpureum* cv. Mott), Guinea grass (*Panicum maximum*), Signal grass (*Bracharia decumbens*)

CONCLUSION

In the results of this study, various grasses had suitable nutritional value and showed that the crude protein of the four grasses showed degradation characteristics: elephant grass (a: 9.88%, b: 64.37%, c: 0.06/h⁻¹), grass mini elephant (a: 16.50%, b: 45.24, and c: 0.05/h⁻¹), guinea grass (a: 7.42%, b: 68.24%, and c: 0.05/h⁻¹), and signal grass reached (a: 6.79%, b: 56.19%, and c: 0.07/h⁻¹). Of the four grasses, it was concluded that they were able to supply the protein requirements for microbial growth in the rumen and the protein needs of the livestock themselves.

DECLARATIONS

Corresponding author

E-mail: ismartoyo@gmail.com

Authors' contribution

Al Wijaya contributes data collection and analysis and the manuscript write up. I Smartoyo and A. Natsir contribute on experiment, idea and research design.

Conflict of interests

The authors have not declared any conflict of interests.

Acknowledgements

This research received funding from Hasanuddin University and the Directorate General of Higher Education, Ministry of Education and Research of the Republic of Indonesia in the Hasanuddin University PDU (Basic Research) 2021 program so that the research can be carried out successfully.

REFERENCES

- Ajayi FT, Babayemi OJ, and Taiwo AA (2007). Effects of *Stylosanthes guianensis* and *Aeschynomene histrix* on the yield, proximate composition and in-situ dry matter and crude protein degradation of *Panicum maximum* (Ntchisi). *Livestock Research for Rural Development*, 19: Article 32. <http://www.lrrd.org/lrrd19/3/ajay19032.htm>
- Akhirany N, Ismartoyo I, Jusoff K, Natsir A, and Ako A (2013). The digestibility, degradation and index value of four local feeds for goat. *World Applied Sciences Journal*, 26: 60–66. <https://doi.org/10.5829/idosi.wasi.2013.26.nrrdsi.26011>
- AOAC (1995). Association of Official Analytical Chemists. Official Method of Analysis. 16th Edition. Washington DC.
- AOAC (2001). Protein (crude) in animal feed, forage (plant tissue), grain, and oilseed. *Journal AOAC International*.
- Baete S and Aregheore EM (2011). Value of *Leucaena leucocephala*, *Gliricidia sepium* and *Ipomea batatas* supplements for apparent nutritional adequacy and maintenance of adult Anglo-Nubian x Fiji local goats on a basal diet of Elephant grass (*Pennisetum purpureum*) in the dry season. *Journal of South Pacific Agriculture*, 15: 10-18. http://repository.usp.ac.fj/9640/1/Wairiu_2011_JOSPA.pdf
- Ballare CL, Scopel AL, and Sanchez RA (1997). Foraging for light: photosensory ecology and agricultural implications. *Plant, Cell and Environment*, 20(6): 820–825. <https://doi.org/10.1046/j.1365-3040.1997.d01-112.x>
- Bamikole MA, Akinsoyinu AO, Ezenwa I, Babayemi OJ, Akinlade J, and Adewumi MK (2004). Effect of six-weekly harvests on the yield, chemical composition and dry matter degradability of *Panicum maximum* and *Stylosanthes hamata* in Nigeria. *Grass and Forage Science*, 59(4): 357–363. <https://doi.org/10.1111/j.1365-2494.2004.00437.x>
- Bozic M, Newton J, Thraen CS, and Gould BW (2012). Mean-reversion in income over feed cost margins: Evidence and implications for managing margin risk by US dairy producers. *Journal of Dairy Science*, 95(12): 7417–7428. <https://doi.org/10.3168/JDS.2012-5818>
- Bonelli E, Zanine A, Souza A, Ferreira D, and Alves G (2013). Ruminal degradability of Guinea grass silage inoculated with *Streptococcus bovis* isolated from bovine rumen combined or not with com wheat bran. *Agricultural Sciences*, 04: 628–634. <https://doi.org/10.4236/as.2013.412084>
- da Silva JKB, da Cunha MV, and dos Santos MVF (2021). Dwarf versus tall elephant grass in sheep feed: which one is the most recommended for cut-and-carry. *Tropical Animal Health and Production*, 53: 93. <https://doi.org/10.1007/s11250-020-02508-y>
- Decandia M, Yiakoulaki M, Pinna G, Cabiddu A, and Molle, G. (2007). Foraging behaviour and intake of goats browsing on mediterranean shrublands. *Dairy Goats, Feeding and Nutrition*, 161–188.
- Ferreira D, Zanine A, Lana R, Divino RM, Alves G, and Mantovani H (2014). Chemical composition and nutrient degradability in elephant grass silage inoculated with *Streptococcus bovis* isolated from the rumen. *Anais Da Academia Brasileira de Ciencias*, 86: 465–473. <https://doi.org/10.1590/0001-37652014112312>
- Guadayo GF, Rayos AA, Merca FE, Tandang AG, Loresco MM, and Angeles AA (2019). Prediction of *In Situ* Ruminal Degradability of Forages in Buffaloes Using the *In Vitro* Gas Production Technique. *Tropical Animal Science Journal*, 42(2): 128–136. <https://doi.org/10.5398/tasj.2019.42.2.128>
- Han X, Chen C, Zhang X, Wei Y, Tang S, Wang J, Tan Z, and Xu, L. (2019). Effects of Dietary Stevioside Supplementation on Feed Intake, Digestion, Ruminal Fermentation, and Blood Metabolites of Goats. *Animals*, 9(2): 32. <https://doi.org/10.3390/ani9020032>
- Hatfield, R. D., & Kalscheur, K. F. (2020). Carbohydrate and Protein Nutritional Chemistry of Forages. *Forages*, 595–607. <https://doi.org/10.1002/9781119436669>
- Ismartoyo (2011). Pengantar teknik penelitian degradasi pakan ternak ruminansia; Introduction to ruminant feed degradation research techniques. *Brilian Internasional*, Sidoarjo, Indonesia
- Jiang B, Yuxiang Z, Tian W, and Fei L. (2020). Nutritive value and ruminal degradation of seven Chinese herbs as forage for Tan sheep. *Bioengineered*, 11(1), 1159–1169. <https://doi.org/10.1080/21655979.2020.1834740>
- Katongole CB, Lumu R, and Lindberg JE (2021). Comparative chemical composition and rumen degradation of common reed and elephant grass in urban/peri-urban dairying systems in Uganda. *Agroecology and Sustainable Food Systems*, 45(6): 892–906. <https://doi.org/10.1080/21683565.2021.1896618>
- Lana RP, Leopoldino WM, Oliveira JS, Veloso RG, Nunes PMM, and Queiroz AC (2007). Parâmetros da degradação protéica ruminal de diferentes alimentos e rações estimados por técnica in vitro. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 59(2): 414–422. <https://doi.org/10.1590/s0102-09352007000200023>
- Larbi A, Smith JW, Kurdi IO, Raji AM and Ladipo DO (1996). Feed value of multipurpose fodder trees and shrubs in West Africa: edible forage production and nutritive value of *Millettia thonningii* and *Albizia lebeck*. *Agroforestry Systems*, 33: 41–50.
- Lemaire G and Belanger G (2020). Allometries in Plants as Drivers of Forage Nutritive Value: A Review. *Agriculture*, 10(1): 5. <https://doi.org/10.3390/agriculture10010005>
- Manaye T, Tolera A, and Zewdu T (2009). Feed intake, digestibility and body weight gain of sheep fed Napier grass mixed with different levels of *Sesbania sesban*. *Livestock Science*, 122(1): 24–29. <https://doi.org/10.1016/j.livsci.2008.07.020>
- Mbuthia EW and Gachui CK (2003). Effect of Inclusion of *Mucuna pruriens* and *Dolichos lablab* Forage in Napier Grass Silage on Silage Quality and on Voluntary Intake and Digestibility in Sheep. *Tropical and Subtropical Agroecosystems*, 1: 123 – 128. <https://www.ccba.uady.mx/publicaciones/journal/2002-2-3/Mbuthia%20and%20Gachui-b.pdf>
- McDonald P, Edwards RA, Greenhalgh JFD, Morgan CA (2002). *Animal Nutrition*. 6th Edition. New York (USA): Ashford Colour Press Ltd.

- Min BR, and Hart SP (2003). Tannins for suppression of internal parasites. *Journal of Animal Science*. 81 (14): E102–E109. https://doi.org/10.2527/2003.8114_suppl_2E102x
- Minson D (1990). *Forage in ruminant nutrition*. Elsevier, San Diego. Pp. 483. <https://www.elsevier.com/books/forage-in-ruminant-nutrition/minson/978-0-12-498310-6>
- Mushandri. 2022. Karakteristik Fermentasi Ransum dalam Rumen Kambing yang Diberi 4 Jenis Hijauan Pakan: Characteristics of Ration Fermentation in the Rumen of Goats Given 4 Types of Forage. Thesis. Hasanuddin University.
- Mutsvangwa T, KL Davies, JJ McKinnon, and DA Christensen (2016) Effects of dietary crude protein and rumen-degradable protein concentrations on urea recycling, nitrogen balance, omasal nutrient flow, and milk production in dairy cows. *Journal of Dairy Science*, 99 (8): 6298-6310. <https://doi.org/10.3168/jds.2016-10917>
- Nichols K, IPC de Carvalho, R Rauch, and J Martín-Tereso (2022) Review: Unlocking the limitations of urea supply in ruminant diets by considering the natural mechanism of endogenous urea secretion, *Animal*, 16 (3): 100537, <https://doi.org/10.1016/j.animal.2022.100537>
- Odedire J. (2013). *In sacco* degradation of *Tephrosia candida* and *Leucaena leucocephala* in mixtures with *Panicum maximum* using fistulated West African dwarf sheep. *Journal of Cell and Animal Biology*, 7: 51–56. <https://doi.org/10.5897/JCAB12.061>
- Ogunwole OA, Akinfemi A, and Akinsoyinu AO (2011). Degradation of crude protein in groundnut cake, guinea grass (*Panicum Maximum*) and rumen epithelial scraping based diets by West African Dwarf sheep. *Nigerian Journal of Animal Production*, 38(1): 106-115. <https://doi.org/10.51791/njap.v38i1.702>
- Ørskov ER, and McDonald I (1979). The estimation of protein degradability in the rumen from incubation measurements weighted according to rate of passage. *The Journal of Agricultural Science*, 92(2), 499–503. <https://doi.org/10.1017/S0021859600063048>
- Ørskov ER (2000). The *in situ* technique for the estimation of forage degradability in ruminants. *Forage evaluation in ruminant nutrition*. CAB International, London. Pp.175-188. <https://doi.org/10.1079/9780851993447.0175>
- Orsoletta AC, Dall AT, Reiter AGV, Kozloski BV, Niderkorn C, and HMN Ribeiro-Filho AD (2017). Associative effects between *Arachis pintoi* and dwarf elephant grass hays on nutritional value in sheep. *Animal Production Science* 58(5) 894-899 <https://doi.org/10.1071/AN15864>
- Patra AK, and Saxena J. (2010). Exploitation of dietary tannins to improve rumen metabolism and ruminant nutrition. *Journal of the Science of Food and Agriculture*, 91(1), 24–37. doi: <https://doi.org/10.1002/jsfa.4152>
- Rasjid, S., and Ismartoyo (2014). Index Value of Goat Feed Based *In sacco* and *In Vivo* Study. *Buletin Nutrisi dan Makanan Ternak*, 10 (1):1-11. <https://journal.unhas.ac.id/index.php/bnmt/article/download/908/579>
- Sari R, WW Jamarun, N Elihasridas, and Yanti G (2021). *In-Vitro* Rument Liquid Characteristics (pH, VFA, and NH₃) From Sugar Cane Top Fermented with Different Levels of *Phanerochaete chrysosporium*. *Proceedings of the International Seminar on Promoting Local Resources for Sustainable Agriculture and Development (ISPLRSAD 2020)*, pp. 190–193.
- Srakaew W, Wachirapakorn C, Cherdthong A, and Wongnen C (2021). Ruminant Degradability and Bypass Nutrients of Alkaline or Steam-Treated Cassava Chip and Corn Grain. *Tropical Animal Science Journal*, 44: 451–461. <https://doi.org/10.5398/tasi.2021.44.4.451>
- Terefe G, Faji M and Mengistu G (2022). Nutritional value and *in situ* degradability of selected forages, browse trees and agro industrial by-products. *Online Journal of Animal and Feed Research*, 12(2): 97-102. DOI: <https://dx.doi.org/10.51227/ojaf.2022.13>
- Sultana S, Khan MJ, Hassan MR, and Khondoker, MAMY (2012). Effects of concentrate supplementation on growth, reproduction and milk yield of Black Bengal goats (*Capra hircus*). *Bangladesh Veterinarian*, 29(1): 7–16. <https://doi.org/10.3329/bvet.v29i1.11884>
- Van Soest PJ (1994). *Nutritional Ecology of the Ruminant*, 2nd ed. Comstock Publishing Associates, Cornell University Press, Ithaca, NY.
- Wang YL, Wang WK, Wu QC, and Yang HJ (2022). The release and catabolism of ferulic acid in plant cell wall by rumen microbes: A review. *Animal Nutrition*, 9: 335–344. <https://doi.org/10.1016/j.aninu.2022.02.003>
- Zulkarnain DR, Ismartoyo, and Harfiah (2014). The Degradation Characteristics of Feed Supplemented *Gliricidia* Leaves in Goat Rumen *in Sacco*. *Jurnal Ilmu dan Teknologi Peternakan*, 3(3): 148-153. <https://journal.unhas.ac.id/index.php/peternakan/article/view/790>