

ESTIMATION OF CRUDE FIBER CONTENT OF A FEED FROM ITS ADF VALUE WHERE THERE IS NO LABORATORY SERVICE

Getachew ASSEFA WOLDEMARIAM 

College of Veterinary Medicine and Animal Sciences, Department of Animal Science, University of Gondar P.O. Box 196, Ethiopia

✉ Email: gawgetachew2@gmail.com

↳ Supporting Information

ABSTRACT: Because of the cost and inaccessibility of laboratory facilities, animal feed formulation at the farm level, in many parts of Ethiopia, is based on feed database information. However, nowadays many laboratories are phasing out the Weende crude fiber (CF) method of analysis. The fiber content of feeds available in most feed databases (including the sub-Saharan Africa feeds composition database) are a result of detergent method analysis (NDF, ADF and lignin). However, CF is still used in poultry feed formulation and forage analysis for horses, in addition to the neutral detergent fiber (NDF) fraction for determining fiber in different countries. Since there is a statistically ($P < 0.01$) difference between the CF and acid detergent fiber (ADF) value of a feed, ADF can't be used directly in place of CF. Therefore, this work aims to formulate a regression equation that could roughly estimate the CF level of a feed from its NDF and ADF values. Considering the strong multicollinearity between NDF and ADF, this study developed separate models for ADF and NDF and compared them based on R^2 and Akaike Information Criterion (AIC), and the ADF-based model provided a better fit. The equations $0.79 \times \text{ADF} - 0.46$, $0.01 + 0.79 \times \text{ADF}$, and $1.37 + 0.62 \times \text{ADF}$ have effectively predicted CF for cereal grains and beans, pulses and byproducts, and also oilseed meals and cakes, respectively. For grass forages, the equation $3.38 + 0.76 \times \text{ADF}$, tested on 10 forages, showed potential but remains unreliable due to its R^2 value below 0.8. Finally, it is concluded that this approach provides a practical alternative for estimating CF where laboratory services or database information are unavailable.

Keywords: Crude Fiber, Estimation, Feed database information, Prediction, Regression.

INTRODUCTION

Dietary carbohydrates can be divided into two basic fractions: fiber and non-fiber carbohydrates (Mirzaei-Aghsaghali and Maheri-Sis, 2011). Fiber is any component in feed that is not digested by mammalian enzymes (Jha and Mishra, 2021). Based on its solubility in water fibers can be grouped into soluble fiber (which dissolves in water) and insoluble fiber. The proximate analysis system developed by the Weende Experiment Station in Germany classified carbohydrates in feed into a more digestible component called nitrogen-free extract (NFE) and a less digestible fibrous component called crude fiber (Singh and Kim, 2021). Crude fiber is a plant cell structural component, including cellulose, hemicelluloses, lignin, and pectin (An et al., 2022; Musa, 2021). The proximate analysis system underestimates the true fiber in the feed. A major problem with this procedure is that the acid and base used in the analysis solubilize some of the true fiber (particularly hemicelluloses, pectin, and lignin), and some cellulose is partially lost too (Musa, 2021). The proximate analysis system only represents a small fraction of the fiber content (average 80% of hemicellulose or pentosans, 50-90% lignin and 50-80% cellulose recovery) (Van Soest and McQueen, 1973). The CF method has a complete recovery of pectins (Möller, 2014).

The other analysis process using neutral and acid detergents by Vax Soest (1963) categorized fiber into neutral detergent fiber (NDF) comprising of cellulose, hemicellulose and lignin, and acid detergent fiber (ADF), largely consisting of cellulose and lignin (Singh and Kim, 2021). The Van Soest detergent fiber system is also affected by unreliability and falls short of accounting for all non-starch polysaccharides (NSP) in the poultry feed ingredients (Singh and Kim, 2021). Non-starch polysaccharides are complex carbohydrates found predominantly in plant cell walls and include components like cellulose, hemicellulose, and pectin. Unlike starch, NSPs cannot be digested by non-ruminants due to their structural complexity and cross-linking, which limits their availability as an energy source.

The Neutral Detergent Fiber (NDF) method has been criticized for not adequately recovering pectin, which is an important part of the cell wall matrix in plants. This omission can lead to an incomplete understanding of the fiber content and its digestibility in poultry diets (Van-Soest et al., 1991). A relatively new feed composition analysis method is Near Infrared Reflectance spectroscopy (NIR). Though NIR method allows rapid and least cost determination of multiple nutrients and characteristics of feeds or forage, it is not available in many places of Ethiopia including the pioneer agricultural university-Haramaya.

RESEARCH ARTICLE
 PII: S222877012500010-15
 Received: December 21, 2024
 Revised: March 15, 2025
 Accepted: March 16, 2025

Because of the cost and inaccessibility of laboratory facilities, animal feed formulation at the farm level, in many parts of Ethiopia, is based on feed database information. However, nowadays many laboratories are phasing out the CF method of analysis. The fiber content of feeds available in most feed databases (including the sub-Saharan Africa feeds composition database) is more of a result of detergent method analysis (NDF, ADF and lignin). However, CF is still used in poultry feed formulation (Singh and Kim, 2021) and forage analysis for horses, in addition to the NDF fraction for determining fiber (Hoffgård, 2022) in different countries. Therefore, the objective of this work is to formulate a regression equation that could roughly estimate the CF level of a feed from its NDF and ADF values.

MATERIALS AND METHODS

Regression models were developed to estimate the CF level of feeds (that could be determined using the Weende proximate analysis system) from their ADF and ADF level using R 4.4.1. The data were taken from Makkar et al. (2024) and INRAE (2024a). First, two candidate models (i.e. using NDF or ADF as predictor variables) were developed for each feed category (i.e. grass forages, cereals grain and bran, pulse seed and byproducts, and oilseed byproducts) and compared using the analysis of variance (ANOVA) in R. Finally, the best-performed model for each feed category was selected for its applicability using a paired sample t-test between the actual and predicted CF values for its non-significance tells us its goodness of fit.

The sample size is determined using Green (1991) and Memon et al. (2020) formula: $n=104+k$; where, n and k are the number of sample size and predictors, respectively. The predator in this case was one (i.e. NDF or ADF) and the minimum sample size would be 105.

The model used in this study was: $Y = \beta_0 + \beta_1\chi_1 + e$; where Y is the response variable (i.e. CF), β_0 is the intercept of the regression line, corresponding to the predicted values when χ_1 (i.e. NDF or ADF) are zero. $\beta_1\chi_1$ is the regression coefficient (β_1) on the independent variables (χ_1 i.e. NDF or ADF). e is the model error (residuals), which defines how much variation is introduced in the model when estimating Y.

RESULTS AND DISCUSSION

Table 1 presents the correlation between CF, NDF, and ADF. The strong correlation between NDF and ADF suggests the presence of high multicollinearity between them indicating they can't be used together in model formulation to predict CF from detergent fiber results. Therefore, separate Models were formulated for ADF and NDF. Choct (2016) noted that, though, the proportion of cellulose and to a lesser extent lignin extracted can be highly variable depending on the ingredient, CF, more or less, represents cellulose and lignin (i.e. ADF) content. However, since there is a significant ($P<0.01$) difference between ADF and CF (Table 1) we can't directly use ADF in place of CF.

Figures 1 and 2 as a partial residual plot showing the linearity assumption of a predictor's (i.e. ADF and NDF) relationship with the dependent variable CF. Both figures show a linear relationship of predator with CF. Where a partial residual plot shows the linear relationship between predictors and dependent variables a linear model can be used (Fox, 2015).

The partial residual plot indicates both the magnitude of the linearity variance and the linearity magnitude and position (Roy et al., 2020).

Table 2 below presents the model parameters and parameters used to compare the models in estimating CF of different categories of feeds from their ADF and NDF contents. In all feed categories, the R^2 value of model one was higher, and the Akaike Information Criterion (AIC) value was lower than model 2. The Delta AIC (ΔAIC) values were much greater than 10. The R^2 value of model one ranges from 0.75 to 0.94. Except for grass forages, the R^2 value of model one of the feed categories was greater than 0.80. The R^2 value equal to 0.8 clearly indicates a very good regression model performance, regardless of the ranges of the ground truth values and their distributions (Chicco et al., 2021). Lower AIC indicates a better fit. If $\Delta AIC >10$, there is strong evidence that the model with the lower AIC is better (Burnham and Anderson, 2002).

Table 1 - Correlation and difference between variables in different feed categories

Parameters	grass forages (Roughages)	Cereal grain and bran	pulse seed and byproducts	Oilseed byproducts
Correlation (NDF and ADF)	0.85	0.86	0.91	0.92
Mean difference between ADF and CF (i.e ADF-CF)	6.14	2.63	4.14	8.61
t-value	-21.97	-11.069	-8.7222	-11.739
SEM	0.74	1.07	1.51	1.81
P-value	< 0.01	< 0.01	< 0.01	< 0.01




Figure 1 - Trends of the relation between crude fiber (CF) and acid detergent fiber (ADF)




Figure 2 - Trends of the relation between crude fiber (CF) and the neutral detergent fiber (NDF).

Table 2 - Estimation of CF of a feed based on its NDF and ADF values.

Models	Model parameters				Model comparison parameters		
	β_0 (Intercept)	β_1 (Coefficient)	R ²	P-value	RSS*	AIC	Δ AIC
grass forages (roughages)							
Model-1: CF~ADF	3.38 ± 1.73	0.76 ± 0.04	0.7454 ^a	<0.001	684.54 ^b	508.24	73.65
Model-2:CF~NDF	-3.90 ± 3.71	0.55 ± 0.05	0.4932 ^b	<0.001	1362.49 ^a	581.89	
SE			0.07		0.60		
P-value			**		**		
Cereals grains and brans							
Model-1: CF~ADF	-0.46 ± 0.27	0.79 ± 0.02	0.9403 ^a	<0.001	297.33 ^b	413.27	184.11
Model-2:CF~NDF	-1.72 ± 0.80	0.35 ± 0.02	0.6555 ^b	<0.001	1716.87 ^a	597.38	
SE			0.10		0.61		
P-value			**		**		
Pulse and their byproducts							
Model-1: CF~ADF	0.01 ± 0.78	0.79 ± 0.03	0.8308 ^a	<0.001	1808.20 ^b	607.50	46.66
Model-2:CF~NDF	-0.65 ± 1.05	0.56 ± 0.03	0.7373 ^b	<0.001	2808.19 ^a	654.16	
SE			0.08		0.91		
P-value			**		**		
Oilseed meals and cake							
Model-1: CF~ADF	1.37 ± 0.89	0.62 ± 0.03	0.8025	<0.001	2282.64 ^b	632.20	72.93
Model-2:CF~NDF	-0.19 ± 1.51	0.45 ± 0.04	0.6070	<0.001	4541.94 ^a	705.13	
SE			0.07		1.11		
P-value			**		**		
RSS: Residual Sum of Squares							

Crude fiber estimation from ADF of grass forages

Table 3 presents the difference between the actual and estimated CF values of 11 grass forages. The paired t-test result indicated no significant difference (P>0.05) between the actual and the predicted values of CF. The predicted CF value of orchard grass was 34.54 and 37.58% of DM. In agreement with this finding, the CF value of Orchard grass was reported to range from 30.2% (Joanna et al., 2007) to 46.45% of DM (Farshadfar, 2012). Glatter et al. (2021) also reported the CF value of meadow hay (35.6%) which is similar to the predicted value of Orchard grass Orchard grass. The 32.94% predicted value of the bamboo leaves reported in this research agrees with the 33.19 % reported by Antwi-Boasiako et al. (2011). Shahowna et al. (2013) reported 54.3 and 47.4 % CF for fresh and fermented sugar cane bagasse, similar to the 48.07% predicted CF value for the same feed using the regression equation developed for grass forages or roughages. The predicted CF value of German grass (33.93%) was within the range of CF value of the German grass at the 3rd (28.30%) and 1st (38.93%) cut reported by Islam et al. (2018). Rahman et al. (2024) also reported the CF value of German grass to be 34.7%. This slight variation in CF content is due to the difference in stage maturity when the sample was analyzed. The predicted CF value (29.14%) for roadside grass was within the range of 32.25 to 36.09 % reported by Haryono et al. (2020) for different cultivars of the roadside grass.

Table 3 - Differences between original CF and model-1 predicted CF for some grass forages*

Feed type/name	Original CF (%)	Literature Source	Predicted CF (%)
Orchardgrass (<i>Dactylis glomerata</i> L): mid-bloom	33.00	Schroeder (2004)	34.54
Orchard grass (<i>Dactylis glomerata</i> L): late-bloom	37.00	"	37.58
Sorghum-Sudan-grass Timothy:			
Late vegetative	27.00	Schroeder (2004)	25.42
Mid-bloom	31.00	"	30.74
Late bloom	31.00	"	45.18
Roadside grass (<i>Stenotaphrum secundatum</i>)	46.27	Selim et al. (2022)	29.14
Banana leaves	29.35	"	31.35
Bamboo leaves	35.56	"	32.94
Sugar cane bagasse	37.89	"	48.07
German grass (<i>Echinochloa polystachya</i>)	39.84	"	33.93
King grass (<i>Pennisetum purpureum</i>)	36.10	Tuturoong et al. (2019)	35.22
Mean	34.91 ^a		34.92 ^a
	SE of the difference	2.59	
Original vs Predicted CF	t-value	-0.0041	
	P-value	0.9968	

* The original crude fiber (CF) used in this table are not used in the model formulation

Crude fiber estimation from ADF of cereals grain and bran

Table 4 presents the differences between the original CF and the CF predicted from ADF using the model for some cereals grain and brans. The insignificant ($P>0.05$) difference observed between the actual and the predicted CF value suggests the good fit of the model in predicting CF from ADF. The CF value of 3.32% predicted from ADF of maize grain was found within the range (2.8 to 4.5%) reported by [Fufa et al. \(2019\)](#) and [Rose and Gupta \(2018\)](#), respectively. The predicted value also falls within the range 2.62 to 3.93 reported by [Radosavljević et al. \(2020\)](#).

[Hossain et al. \(2008\)](#) reported 11.38% CF for wheat bran, similar to the 11.64% CF predicted from the ADF of wheat bran. In agreement with this [Liu et al. \(2024\)](#) also reported 10.94% of CF for wheat bran. The CF values of 3.2% and 5.23% predicted from ADF of wheat and barley grain, respectively, were similar to 3.0% and 5.23% reported for wheat and barley grain by [Hossain et al. \(2008\)](#) and [Venslovas et al. \(2024\)](#), respectively. The CF values of 5.81, 5.44, and 7.05% predicted from ADF of Sorghum grain were similar to 5.90, 5.40, and 6.50 % reported by [Kumar et al. \(2019\)](#) for different sorghum varieties. In agreement with this finding [Treviño-Salinas et al. \(2021\)](#) also reported 6.07 to 9.09 % of CF for different varieties of sorghum grains. The CF value of 4.17% predicted from the ADF of Sorghum HB2 was similar to the 4.17% reported by [Banna and Arifuddin \(2024\)](#).

Table 4 - Differences between original CF and model-1 predicted CF for some cereals grain and bran *

Feed type/name	ADF (%)	Original CF (%)	Literature source	Predicted CF (%)
Maize grain	4.79	1.10	Jaishankar et al. (2021)	3.32
Wheat bran	15.32	14.07	Ning et al. (2022)	11.64
Wheat	4.63	2.71	"	3.20
Barley grains	7.20	9.0	Asma et al. (2021)	5.23
Corn	3.96	1.82	Sheikhhasan et al. (2020)	2.67
Sorghum HB1	7.94	1.96	Salinas et al. (2006)	5.81
Sorghum HB2	5.86	2.67	"	4.17
Sorghum HB3	9.51	4.17	"	7.05
Sorghum HB4	24.24	9.02	"	18.69
Sorghum HB5	25.47	6.80	"	19.66
Sorghum HB7	7.47	1.71	"	5.44
Mean		5.00 ^a		7.90 ^a
	SE of the difference		2.23	
Original vs Predicted CF	t-value		-1.9943	
	P-value		0.07409	

* The ADF and original CF used in this table are not used in model formulation. CF: crude fiber;; ADF: acid detergent fiber

Crude fiber estimation from ADF of pulse seeds and byproducts

Table 5 presents the differences between the original CF and the CF predicted from ADF using the model formulated for pulse seeds and byproducts. The insignificant ($P>0.05$) difference observed between the original and the predicted CF value suggests the ability of the model to predict CF from ADF. The CF values of 27.98%, 32.48%, and 9.16% predicted from ADF of the cowpea haulms, cowpea Pod husks, and faba bean seeds, respectively, were similar to 27.5%, 31.8 %, and 8.9 % reported by [Antwi et al. \(2014\)](#), [Abebe and Alemayehu \(2022\)](#) and [Micek et al. \(2015\)](#), respectively for similar ingredients. The CF value of 5.15% predicted from ADF of cowpea seed was also comparable to the 5.66% reported by [Gutema and Tolesa \(2024\)](#). The CF value of 11.31% predicted from ADF of Lupin seed was found within the range (10.0 to 16.0%) reported by [Abraham et al. \(2019\)](#). [Uzun and Okur \(2023\)](#) also reported 11.75 % CF for Blue lupin. The CF values of 5.15% and 7.36% predicted values from ADF of the Adzuki bean and pigeon pea were similar to the 4.71 ± 0.54 % and 6.6% reported by [Sai-Ut et al. \(2010\)](#) and [Saxena et al. \(2010\)](#), respectively. The CF values of 4.32%, 7.81% and 5.90% predicted from ADF of Chickpea, Faba bean, and Common vetch were comparable to the 3.9 %, 7.72 %, and 3.80–7.17 % reported by [INRAE \(2024b\)](#), [Smit et al. \(2021\)](#), and [Huang et al. \(2017\)](#), respectively.

Crude fiber estimation from ADF of Oilseed byproducts

Table 6 presents the differences between the original CF and the CF predicted from ADF using the model formulated for oilseed byproducts. The insignificant ($P>0.05$) difference between the original and the predicted CF suggests the ability of the model to predict CF from ADF. The CF values of 8.31%, and 4.64% predicted from ADF of rapeseed cake and Soybean meal were similar to the 7.9 % and 4.40% reported by the [National Dairy Development Board \(2012\)](#) of India and [Makkar et al. \(2024\)](#), respectively. In agreement with this finding [Tang et al. \(2024\)](#) also reported 8.26 % CF for rapeseed cake.

The CF value of 10.93% predicted from ADF of canola meal was in the range 8.97 to 11.4 % reported by [Birmani et al. \(2019\)](#). [Kaiser et al. \(2022\)](#) also reported 10.1% CF of the canola meal. The CF value of 17.92% predicted from ADF of Noug seed cake was similar to the 17.65 % reported by [Amare et al. \(2021\)](#). The CF value of 17.12% predicted from ADF of Sesame meal was less than the 9.86 % reported by [Elfaki and Unal \(2023\)](#). The Difference is due to their ADF content. The former had 25.41% ADF, while the latter had 13.83%. The CF value of 15.11% predicted from ADF of Sunflower cake was within the range (11.6-23.89%) reported by [Swain et al. \(2023\)](#). The CF values of 8.44% and 4.68% predicted from the ADF of flaxseed cake and soybean meal were similar to the 8.8 % and 5.44 % reported by [Nehmeh et al. \(2022\)](#) and [Etiosa et al. \(2018\)](#), respectively. [Dunmire et al. \(2021\)](#) also reported CF values ranging from 4.27 to 5.17% for soybean meal produced from different varieties of soybean, which is similar to the above-mentioned predicted CF value for soybean meal.

Table 5 - Original CF and model-1 predicted CF for some pulse seeds and byproducts *

Feed type/name	NDF (%)	ADF (%)	Original CF (%)	Literature source	Predicted CF (%)
Cowpea haulms	49	35.4	29.9	Li et al. (2021)	27.98
Cowpea Pod husks	54.2	41.1	31.8	"	32.48
Faba bean seeds	22.06	11.58	9.72	Meng et al. (2021)	9.16
cowpea seed	16.6	6.5	5.6	Makkar et al 2024	5.15
Lupins (al bus)	17.2	14.3	10.27	Sipas et al. (1997)	11.31
Adzuki bean	12.70	6.50	4.76	"	5.15
Pigeon pea	13.70	9.30	8.07	"	7.36
Chickpea Kabuli	11.90	5.46	2.93	"	4.32
Faba bean	12.79	9.87	8.41	"	7.81
Common vetch	21.90	7.46	5.10	"	5.90
		Mean	11.656 ^a		11.662 ^a
Original vs Predicted CF		SE of the difference		4.5733	
		t-value		-0.012801	
		P-value		0.9901	

* The NDF, ADF, and original CF used in this table are not used in the model formulation. CF: crude fiber; NDF: neutral detergent fiber; ADF: acid detergent fiber

Table 6 - Original CF and model-1 predicted CF for some Oilseed byproducts *

Feed type/name	NDF (%)	ADF (%)	Original CF (%)	Literature sources	Predicted CF (%)
Rapeseed cake	17.80	26.03	11.19	Renata et al. (2018)	8.31
Soybean meal	8.21	5.28	3.89	Tanawong (2013)	4.64
Canola meal	22.64	15.42	10.50	"	10.93
Noug seed cake	34.5	26.7	22.0	Moges et al. (2016)	17.92
Sesame seed meal	37.50	24.25	9.00	Mahmoud and Wafaa (2014)	16.41
Cottonseed Cake	61.53	17.70	12.10	Idrissou et al. (2020)	12.34
Sesame meal	39.35	25.41	3.28	Omer et al. (2019)	17.12
Flaxseed cake	14.2	11.4	8.1	Niyonshuti and Kirkpinar (2024)	8.44
Sunflower cake	32.59	36.56	22.16	Renata et al. (2018)	15.11
Soybean	9.36	13.99	5.34	"	4.68
		Mean	10.76		11.59
Original CF vs Predicted CF		SE of the difference		2.6353	
		t-value		-1.6653	
		P-value		0.1198	

CF: crude fiber; NDF: neutral detergent fiber; ADF: acid detergent fiber

CONCLUSION

Where there is no laboratory service and the database information for crude fiber values for a feed in question, it is possible to estimate the CF value from the ADF value of a feed. Since there is a statistically significant ($P < 0.01$) difference between the CF and ADF value of a feed, ADF can't be used directly in place of CF. The regression equation $0.79 \times \text{ADF} - 0.46$, $0.01 + 0.79 \times \text{ADF}$, and $1.37 + 0.62 \times \text{ADF}$ can be used for cereals grains and brans, pulse and their byproducts, and oilseed meals and cake, respectively. However, for the R^2 value less than 0.8, the regression equation $3.38 + 0.76 \times \text{ADF}$ formulated for grass forages, even though tested on 10 forages and found effective, the model is not reliable.

DECLARATIONS

Corresponding author

Correspondence and requests for materials should be addressed to Getachew Assefa Woldemariam; E-mail: gawgetachew2@gmail.com; ORCID: <https://orcid.org/0000-0002-0593-0739>

Supplementary Information

The supplementary materials are available at <https://ojafr.com/> (Volume 15, Issue 2; Pages 79-88).

Author's Contribution

I contribute to data analysis, interpretation, discussion, and the write-up of the manuscript.

Competing interests

The author has not declared any competing interests.

Consent to publish

The author has reviewed and approved the final manuscript for publication.

Funding

The research has no founding resources.

Acknowledgments

I thank the University of Gondar, for the uninterrupted facilities at the university, home, and the office for conducting and completing the study. My heartfelt gratitude also goes to the Research Ethics Review Committee of the College of Veterinary Medicine and Animal Sciences for their meticulous review and approval.

REFERENCES

- Abebe BK, and Alemayehu MT (2022). A review of the nutritional use of cowpea (*Vigna unguiculata* L. Walp) for human and animal diets. *Journal of Agriculture and Food Research*, 10: 100383. DOI: <https://doi.org/10.1016/j.jafr.2022.100383>.
- Abraham EM, Ganopoulos I, Madesis P, Mavromatis A, Mylona P, Nianiou-Obeidat I, et al. (2019). The use of lupin as a source of protein in animal feeding: Genomic tools and breeding approaches. *International Journal of Molecular Sciences*, 20(4): 851. DOI: <https://doi.org/10.3390/ijms20040851>.
- An Y, Lu W, Li W, Pan L, Lu M, Cesarino I, et al. (2022). Dietary fiber in plant cell walls—the healthy carbohydrates. *Food Quality and Safety*, 6: fyab037. <https://doi.org/10.1093/fqsafe/fyab037>
- Antwi-Boasiako C, Coffie GY, and Darkwa NA (2011). Proximate composition of the leaves of *Bambusa ventricosa*, *Oxytenanthera abyssinica* and two varieties of *Bambusa vulgaris*. *Scientific Research and Essays*. 6(34): 6835-6839. DOI: <https://doi.org/10.5897/sre11.797>.
- Amare SS, Habtamu AK and Yohannes DA (2021). Effects of replacement noug seed cake (*Guizotia abyssinica*) with treated tagasaste (*Chamaecytisus palmensis*) leaf meal on egg quality traits of layer hens. *International Journal of Agricultural Science and Food Technology*, 7(3): 277-284. DOI: <https://dx.doi.org/10.17352/2455-815X.000120>.
- Antwi C, Osafo ELK, Donkoh A, and Adu-Dapaah H (2014). Chemical composition, gas production and degradation characteristics of haulms of improved dual-purpose Cowpea (*Vigna unguiculata* L. walp) cultivars. *Livestock Research for Rural Development* 26 (11): Article #194. <http://www.lrrd.org/lrrd26/11/antw26194.html>.
- Asma A, Imen C, Roua R, and Ayed MH (2021). Effect of partial dietary substitution of Carob (*Ceratonia siliqua* L.) to barley grains on diet digestibility in growing rabbits. *Journal of New Sciences, Agriculture and Biotechnology*, 79 (1): 4580-4585. <https://www.jnsiences.org/agri-biotech/114-volume-79/651>.
- Banna MZA, and Arifuddin W (2024). Characterizing agronomic, morphological, nutritional, and phytochemical traits of local sorghum (*Sorghum bicolor* L.) in Indonesia. *Jurnal Ilmiah Pertanian*, 21(2). DOI: <https://doi.org/10.31849/jip.v21i2.16672>.

- Birman MW, Nawab A, Ghani MW, Gaowen, Liu W, Xiao M, et al. (2019). Canola meal in poultry nutrition; solubility test to determine its protein quality. International Journal of Life Sciences Research, 3: 67-69. <https://www.researchpublish.com/upload/book/Canola%20meal%20in%20poultry%20nutrition-7811.pdf>
- Burnham KP, and Anderson DR (2002). Model Selection and Multimodel Inference: A Practical Information-Theoretic Approach (2nd ed.). Springer-Verlag, New York. <https://doi.org/10.1007/b97636>
- Chicco D, Warrens MJ, and Jurman G (2021). The coefficient of determination R-squared is more informative than SMAPE, MAE, MAPE, MSE and RMSE in regression analysis evaluation. Peer J Computer Science, 7: e623. DOI: <https://doi.org/10.7717/peerj-cs.623>.
- Choct M (2016). Measurements of nutrients and nutritive value In M. R. Bedford, M. Choct, and H. V. M. O'Neill (Ed.), Nutrition experiments in pigs and poultry: A practical guide. CAB International, pp. 74-98. DOI: <https://doi.org/10.1079/9781780647005.0074>.
- Dunmire KM, Braun MB, Evans CE, Fallen BD, Stark CR, and Paul CB (2021). Determining Amino Acid Digestibility of Soybean Meal from South Carolina Soybean Varieties Fed to Broilers. Kansas Agricultural Experiment Station Research Reports. 7(10):21. <https://newprairiepress.org/kaesrr/vol7/iss10/21/>.
- Elfaki MOA, and Unal K (2023). Determination of feed value, in vitro digestibility and enteric methane production of black seed (*nigella sativa*) meal. Journal of Scientific and Engineering Research, 10(6): 173-177. <https://www.researchgate.net/publication/374229098>.
- Etiosa O, Chika N, and Benedicta A (2018). Mineral and proximate composition of soya bean. Asian Journal of Physical and Chemical Sciences, 4(3): 1-6. DOI: <https://doi.org/10.9734/ajopacs/2017/38530>.
- Farshadfar M (2012). Evaluation of the forage yield and quality in some accessions of *Dactylis Glomerata* under irrigated conditions. Annals of Biological Research, 3(4): 1694-1699. <https://www.scholarsresearchlibrary.com/articles/evaluation-of-the-forage-yield-and-quality-in-some-accessions-of-dactylis-glomerata-under-irrigated-conditions.pdf>
- Fox J (2015). Applied regression analysis and generalized linear models (3rd ed.). Thousand Oaks, SAGE Publications, CA. <https://dokumen.pub/applied-regression-analysis-and-generalized-linear-models-3nbsped-1452205663-9781452205663.html>.
- Fufa N, Aberra S, and Girma D (2019). Effect of storage structures and storage period on grain quality of maize (*Zea mays* L.): the case of West Shawa Zone, Bako, Ethiopia. International Journal of Current Research and Academic Review, 7(4): 12-19. DOI: <https://doi.org/10.20546/ijcrar.2019.704.003>.
- Glatter M, Bochnia M, Wensch-Dorendorf M, Greef JM, and Zeyner A (2021). Feed intake parameters of horses fed soaked or steamed hay and hygienic quality of hay stored following treatment. Animals, 11 (2729): 1-15. DOI: <https://doi.org/10.3390/ani11092729>.
- Green SB (1991). How many subjects does it take to do a regression analysis. Multivariate Behavioral Research, 26(3): 499-510. DOI: https://doi.org/10.1207/s15327906mbr2603_7.
- Gutema T, and Tolesa GN (2024). Effects of traditional processing techniques on nutritional quality and sensory acceptability of value-added products made from cowpea (*Vigna unguiculata* L. walp.) produced in Ethiopia. The North African Journal of Food and Nutrition Research, 8(17): 32-43. DOI: <https://doi.org/10.51745/najfnr.8.17.32-43>.
- Haryono E, Umami N, Ananta D, Bachruddin Z, Suhartanto B, Hanim et al. (2020). Nutrient content, fiber fraction and ethanol production of three cultivars (*Pennisetum purpureum* Scumach.). E3S Web of Conferences, 200: 03008. DOI: <https://doi.org/10.1051/e3sconf/202020003008>.
- Hoffgård M (2022). The effect of harvest time of timothy on digestibility in equine with in sacco and in vitro methods. Norwegian University of Life Science. <https://nmbu.bragu.unit.no/nmbu-xmlui/handle/11250/3038226>
- Hossain FMA, Khan RI, Bostami ABMR, Mondal A, and Azad MAK (2008). Replacement of Wheat Bran by Biogas Slurry and their Effect on Intake and Digestibility of Deshi Sheep. Journal of Agroforestry and Environment, 2(1): 147-150. <https://jagroforenviron.com/wp-content/uploads/2018/09/30-Replacement-of-wheat-bran-by-biogas-slurry-and-their-effect-on-intake-and-digestibility-of-deshi-sheep-Hossain.pdf>.
- Huang YF, Gao XL, Nan ZB, and Zhang ZX (2017). Potential value of the common vetch (*Vicia sativa* L.) as an animal feedstuff: a review. Journal of Animal Physiology and Animal Nutrition, 101(5): 807-823. DOI: <https://doi.org/10.1111/jpn.12617>.
- Idrissou Y, Worogo HSS, Assani AS, Ayena JA., Assogba BGC, and Traoré IA (2020). Cottonseed cake replacement by soybean pulp in the diet of West African Dwarf lambs in Benin: zootechnical and economic performances. Journal of Livestock and Veterinary Medicine in Tropical Countries 73(2): 107-111. DOI: <https://doi.org/10.19182/remvt.31875>.
- INRAE, CIRAD and AFZ (2024a). Feed tables: Composition and nutritive values of feed materials for ruminants, pigs, poultry, rabbits, horses, and salmonids. Retrieved [10/12/2024], from <https://www.feedtables.com/content/table-dry-matter>
- INRAE, CIRAD, and AFZ (2024b). Table of dry matter. Retrieved 15 Dec 2024, from <https://www.feedtables.com/content/table-dry-matter>.
- Islam MM, Khan MMH, Uddin MJ, and Islam MJ (2018). Biomass and nutritional evaluation of German grass (*Echinochloa polystachya*) cultivated on floating bed. Bangladesh Journal of Animal Science, 46(4): 249-257. DOI: <https://doi.org/10.3329/bjas.v46i4.36966>.
- Jaishankar N, Ramachandra B, and Thirumalesh T (2021). Effect of urea ammoniated sugarcane trash based diets on growth potentiality of Nari Suwana x Kenguri F1 lamb. The Pharma Innovation Journal, 10(11): 138-142. <https://www.thepharmajournal.com/archives/2021/vol10issue11S/PartC/S-10-10-99-864.pdf>
- Jha R, and Mishra P (2021). Dietary fiber in poultry nutrition and their effects on nutrient utilization, performance, gut health, and on the environment: a review. Journal of Animal Science and Biotechnology, 12(1): 51. DOI: <https://doi.org/10.1186/s40104-021-00576-0>.
- Joanna J, Jolanta J, and Grazyna C (2007). Fodder quality from orchard grass and its mixtures with the legumes in the aspect of different nitrogen fertilization. Zbornik radova, 44: 2007. <https://scindeks-clanci.ceon.rs/data/pdf/0351-4781/2007/0351-47810702143J.pdf>.
- Kaiser F, Harbach H and Schulz C (2022). Rapeseed proteins as fishmeal alternatives: A review. Reviews in Aquaculture, 14(4): 1887-1911. DOI: <https://doi.org/10.1111/raq.12678>
- Kumar A, Kewat R, Yadav V, Kumar D, and Lal K. (2019). Biochemical evaluation of different sorghum genotypes for high mineral and crude fiber content. International Journal of Chemical Studies, 7(5): 1452-1454. <https://www.chemjournal.com/archives/2019/vol7issue5/PartZ/7-5-191-727.pdf>.
- Li H, Yin J, Tan B, Chen J, Zhang H, Li Z, et al. (2021). Physiological function and application of dietary fiber in pig nutrition: A review. Animal Nutrition, 7(2): 259-267. DOI: <https://doi.org/10.1016/j.aninu.2020.11.011>
- Liu X, Yun X, Cheng Z, Guo Y, Yuan J, and Nie W. (2024). Determination and prediction of standardized ileal amino acid digestibility of wheat bran in broiler chickens. Animal Bioscience, 37 (10): 1788-1798. <https://doi.org/10.5713/ab.24.0003>.

- Mahmoud AEM, and Wafaa MAG (2014). Effect of partial substitution of dietary protein by nigella sativa meal and sesame seed meal on performance of Egyptian lactating buffalo. *Asian Journal of animal and Veterinary Advances* 9(8): 489-498. DOI: <https://scialert.net/abstract/?doi=ajava.2014.489.498>.
- Makkar HPS, Tran G, Heuzé V, and Ankers P (2024). Feedipedia: Soybean meal. Animal Feed Resources Information System. <https://www.feedipedia.org/node/674>.
- Memon MA, Ting H, Cheah J, Thurasamy R, Chuah F, and Cham TH (2020). Sample Size for Survey Research: Review and Recommendations. *Journal of Applied Structural Equation Modeling*, 4(2): 1-20. https://jasemjournal.com/wp-content/uploads/2020/08/Memon-et-al_JASEM_Editorial_V4_Iss2_June2020.pdf.
- Meng Z, Liu Q, Zhang Y, Chen J, Sun Z, Ren C, et al. (2021). Nutritive value of faba bean (*Vicia faba* L.) as a feedstuff resource in livestock nutrition: A review. *Food Science & Nutrition*, 9(9): 5244-5262. DOI: <https://doi.org/10.1002/fsn3.2342>.
- Micek P, Kowalski ZM, Kulig B, and Kański J (2015). Effect of variety and plant protection method on chemical composition and in vitro digestibility of Faba bean (*Vicia faba*) seeds. *Annals of Animal Science*, 15(1): 143-154. DOI: <https://doi.org/10.2478/aoas-2014-0080>.
- Mirzaei-Aghsaghali A and Maheri-Sis N. (2011). Importance of “physically effective fiber” in ruminant nutrition: A review. *Annals of Biological Research*, 2(3): 262-270. <https://www.scholarsresearchlibrary.com/articles/importance-of-physically-effective-fibre-in-ruminant-nutrition-a-review.pdf>
- Moges D, Mengistu U, Getachew A, Mohammed YK, and Tilahun S (2016). Effect of concentrate supplementation to free ranging dromedary camels on yield, physicochemical quality and fatty acid profile of milk. *Livestock Research for Rural Development*, 28(6): Article #99. <https://lrrd.cipav.org.co/lrrd28/6/dere28099.html>.
- Möller J (2014). Comparing methods for fibre determination in food and feed. A White Paper from FOSS. https://www.fossanalytics.com/-/media/files/documents/papers/laboratories-segment/fibre-methods-compared_gb.pdf.
- Musa AMT (2021). Chemical composition and in vitro digestibility of some range plants. *Open Journal of Plant Science*, 6(1): 094-098. <https://dx.doi.org/10.17352/ojps.000040>
- National Dairy Development Board (2012). Nutritive value of commonly available feeds and fodders in India. Animal Nutrition booklet. Retrieved from <https://www.nddb.coop/sites/default/files/pdfs/Animal-Nutrition-booklet.pdf>.
- Nehmeh M, Rodriguez-Donis I, Cavaco-Soares A, Evon P, Gerbaud V, and Thiebaut-Roux S (2022). Bio-refinery of oilseeds: oil extraction, secondary metabolites separation towards protein meal valorisation—A Review. *Processes*, 10(5): 841. DOI: <https://doi.org/10.3390/pr10050841>.
- Ning R, Cheng Z, Liu X, Ban Z, Guo Y, and Nie W (2022). Evaluating and predicting net energy value of wheat and wheat bran for broiler chickens. *Animal Bioscience*, 35(11): 1760-1770. DOI: <https://doi.org/10.5713/ab.21.0501>.
- Niyonshuti E, and Kirkpinar F (2024). Comparison of nutrient composition and potential feed value of different cakes obtained by cold pressed method. *Journal of the Hellenic Veterinary Medical Society*, 75(3): 8017-8026. DOI: <https://doi.org/10.12681/jhvms.36360>.
- Omer HAA, Ahmed SM, Abdel-Magid SS, Bakry BA, El-Karamany MF, and El-Sabaawy EH (2019). Nutritional impact of partial or complete replacement of soybean meal by sesame (*Sesamum indicum*) meal in lambs rations. *Bulletin of the National Research Centre*, 43(1): 98. DOI: <https://doi.org/10.1186/s42269-019-0140-8>.
- Radosavljević M, Milašinović-šeremešić M, Terzić D, Jovanović Ž, Srdić J, and Nikolić V (2020). Grain chemical composition of dents, popping maize and sweet maize genotypes. *Journal on Processing and Energy in Agriculture*, 24(2): 77-80. DOI: <https://doi.org/10.5937/jpea24-28790>.
- Rahman MA, Redoy MRA, Shuvo AA, Chowdhury R, Hossain E, Sayem SM, et al.(2024). Influence of herbal supplementation on nutrient digestibility, blood biomarkers, milk yield, and quality in tropical crossbred cows. *PLoS ONE* 19(11): e0313419. <https://doi.org/10.1371/journal.pone.0313419>
- Renata K, Luba B, Patrícia V, Branislav V, Martin G, Daniel B, et al. (2018). Fiber content and in vitro digestibility of oilseeds and their cakes. In: O. Stastnik and E. Mrkvicova (ed.) *NutriNET 2018*, pp. 53-59. https://nutrinet.mendelu.cz/wcd/w-rek-nutrinet/2018/nutrinet_2018_final.pdf#page=54.
- Rose A, and Gupta A (2018). Comparative analysis and optimization by utilizing different maize varieties (*Zea mays* L.). *Journal of Pharmacognosy and Phytochemistry*, 7(6): 1878-1882. <https://www.phytojournal.com/archives/2018/vol7issue6/PartAG/7-6-4-385.pdf>.
- Roy P, Pal SC, Arabameri A, Chakraborty R, Pradhan B, Chowdhuri I, et al. (2020). Novel ensemble of multivariate adaptive regression spline with spatial logistic regression and boosted regression tree for gully erosion susceptibility. *Remote Sensing*, 12(3284):1-35. DOI: <https://doi.org/10.3390/rs12203284>
- Sai-Ut S, Ketnawa S, Chaiwut P, and Rawdkuen S (2010). Biochemical and functional properties of proteins from red kidney, navy and adzuki beans. *Asian Journal of Food and Agro-Industry*, 2(04): 493-504. https://www.academia.edu/82750105/Biochemical_and_functional_properties_of_proteins_from_red_kidney_navy_and_adzuki_beans.
- Salinas I, Pró A, Salinas Y, Sosa E, Becerril CM, Cuca M, et al. (2006). Compositional variation amongst sorghum hybrids: Effect of kafirin concentration on metabolizable energy. *Journal of Cereal Science*, 44(3): 342-346. DOI: <https://doi.org/10.1016/j.jcs.2006.08.008>.
- Saxena KB, Kumar RV, and Sultana R (2010). Quality nutrition through pigeonpea—a review. *Health*, 02(11):1335-1344. DOI: <https://doi.org/10.4236/health.2010.211199>.
- Schroeder JW (2004). Forage nutrition for ruminants. North Dakota State University Fargo, North Dakota. <https://www.library.nd.gov/statedocs/NDSUExtensionService/as125020090702.pdf>
- Selim ASM, Hasan MN, Rahman MA, Rahman MM, Islam MR, Bostami A, et al. (2022). Nutrient content and in vitro degradation study of some unconventional feed resources of Bangladesh. *Heliyon*, 8(5): e09496. DOI: <https://doi.org/10.1016/j.heliyon.2022.e09496>.
- Shahowna EM, Mahala AG, Mokhtar AM, Amasaib EO, and Attaelmnan B (2013). Evaluation of nutritive value of sugar cane bagasse fermented with poultry litter as animal feed. *African Journal of Food Science and Technology*, 4(5): 106-109. <https://www.researchgate.net/publication/280320450>.
- Sheikhhasan BS, Moravej H, Ghaziani F, Esteve-Garcia E, and Kim WK (2020). Relationship between chemical composition and standardized ileal digestible amino acid contents of corn grain in broiler chickens. *Poultry Science*, 99(9), 4496-4504. DOI: <https://doi.org/10.1016/j.psj.2020.06.013>.

- Singh AK, and Kim WK (2021). Effects of dietary fiber on nutrients utilization and gut health of poultry: A Review of challenges and opportunities. *Animals (Basel)*, 11(1):181. DOI: <https://doi.org/10.3390/ani11010181>.
- Sipas S, Mackintosh JB, and Petterson DS (1997). The chemical composition and nutritive value of Australian pulses. Grains Research and Development Corporation, Kingston, Australia. pp. 5-62. <https://library.dpi.nsw.gov.au/books/20/>.
- Smit MN, He L, and Beltranena E (2021). Feeding different cultivars and quality levels of faba bean to broiler chickens. *Translational Animal Science*, 5(3), txab094. DOI: <https://doi.org/10.1093/tas/txab094>.
- Swain KB, Beura CK, Naik PK, Kumar D, Mishara SK (2023). Alternative feed resources for sustainable poultry production. In: Srivastava A.K., Shukala P.K., Amitav Bhattacharyya and Singh M.K. (editors), Recent trends in sustainable poultry production. Satish Serial Publishing House, Delhi, India. P. 105 -124. <https://krishi.icar.gov.in/jspui/bitstream/123456789/75168/1/book%20chapteripsacon2022book.pdf>
- Tanawong M (2013). Evaluation of the nutritional value of Canola meal, 00-rapeseed meal and 00-rapeseed expellers fed to pigs. Dissertation, University of Illinois. Urbana-Champaign. <https://nutrition.ansci.illinois.edu/sites/default/files/DissertationMaison.pdf>
- Tang H, Fengn G, Zhao J, Ouyang Q, Liu X, Jiang X, et al. (2024). Determination and prediction of amino acid digestibility in rapeseed cake for growing-finishing pigs. *Animals*, 14: 2764. <https://doi.org/10.3390/ani14192764>
- Treviño-Salinas M, Perales-Torres A, Castillo-Ruiz O, Montes-García N, Lizarazo-Ortega C, Navarro-Cortez R, et al. (2021). Proximal analysis and profile of fatty acids on six varieties of white grain sorghum with potential use in human consumption. *CYTA – journal of food*, 19(1): 547–551. <https://doi.org/10.1080/19476337.2021.1928757>.
- Tuturoong RAV, Malalantang SS, and Moningkey SAE (2019). Assessment of the nutritive value of corn stover and king grass in complete feed on Ongole steer calves' productivity. *Veterinary World*, 13(4): 801-806. DOI: <https://doi.org/10.14202/vetworld.2020.801-806>.
- Uzun T and Okur AA (2023). Impacts of different processes on the nutritional and antinutritional contents of white and blue lupin seeds and usage possibilities for sustainable poultry production. *Animals* 13: 3496. DOI: <https://doi.org/10.3390/ani13223496>.
- Vax Soest PJ (1963). The use of detergents in the analysis of fibrous feeds: ii. a rapid method for the determination of fiber and lignin. *Journal of Association of Official Agricultural Chemists*, 46(5): 829-835. DOI: <https://doi.org/10.1093/jaoac/46.5.829>.
- Van-Soest PJ, Robertson JB, and Lewis BA (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal of Dairy Science*, 74: 3583-3597. [https://www.journalofdairyscience.org/article/S0022-0302\(91\)78551-2/pdf](https://www.journalofdairyscience.org/article/S0022-0302(91)78551-2/pdf)
- Van Soest PJ, and McQueen RW (1973). The chemistry and estimation of fibre. *Proceedings of the Nutrition Society*, 32(3): 123-130. DOI: <https://doi.org/10.1079/pns19730029>.
- Venslovas E, Kochiieru Y, Janaviciene S, Merkevicute-Venslove L, Almgodad M, Bartkevics V, et al. (2024). impact of harvest delay and barley variety on grain nutritional composition and mycotoxin contamination. *Journal of Fungi*, 10(11): 738. DOI: <https://doi.org/10.3390/jof10110738>.

Publisher's note: Scienceline Publication Ltd. remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access: This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <https://creativecommons.org/licenses/by/4.0/>.